Package 'BiTSLS'

May 28, 2025

Type Package
Title Bidirectional Two-Stage Least Squares Estimation
Version 0.1.0
Date 2025-05-26
Description Implements bidirectional two-stage least squares (Bi-TSLS) estimation for identifying bidirectional causal effects between two variables in the presence of unmeasured confounding. The method uses proxy variables (negative control exposure and outcome) along with at least one covariate to handle confounding.
License MIT + file LICENSE
Encoding UTF-8
Imports stats
RoxygenNote 7.3.1
Depends R (>= 3.5.0)
NeedsCompilation no
Author Jiaqi Min [aut, cre]
Maintainer Jiaqi Min <fhoneysuckle7@gmail.com></fhoneysuckle7@gmail.com>
Repository CRAN
Date/Publication 2025-05-28 15:30:06 UTC

Contents

	Bi_TSLS	 	• •	 • •	 • •	•	 • •	 	 •	•••	 •		 • •	•	 2
Index															3

Bi_TSLS

Description

Performs bidirectional two-stage least squares (Bi-TSLS) estimation to identify bidirectional causal effects between X and Y using proxy variables Z and W. Requires at least one covariate in addition to X, Y, Z, and W.

Usage

 $Bi_TSLS(data, R_w = 0, R_z = 0)$

Arguments

data	A data frame containing at least these variables:
	• X: Treatment/exposure variable
	• Y: Outcome variable
	• Z: Negative control exposure (proxy variable)
	• W: Negative control outcome (proxy variable)
	• At least one additional covariate (required)
R_w	Sensitivity parameter for W (default = 0)
R_z	Sensitivity parameter for Z (default = 0)

Value

A list containing:

- beta_XY: Estimated effect of X on Y
- beta_YX: Estimated effect of Y on X

Examples

```
# Create example data with covariate
n <- 1000
data <- data.frame(
  X = rnorm(n),
  Y = rnorm(n),
  Z = rnorm(n),
  W = rnorm(n),
  V = rnorm(n) # At least one covariate is required
)
# Run BiTSLS
result <- Bi_TSLS(data)
print(result)
```

Index

Bi_TSLS, 2