

# Package ‘LinearRegressionMDE’

January 20, 2025

**Title** Minimum Distance Estimation in Linear Regression Model

**Version** 1.0

**Description** Consider linear regression model  $Y = Xb + \text{error}$  where the distribution function of errors is unknown, but errors are independent and symmetrically distributed. The package contains a function named LRMDE which takes Y and X as input and returns minimum distance estimator of parameter b in the model.

**Depends** R (>= 3.2.2)

**License** GPL-2

**LazyData** true

**NeedsCompilation** no

**Author** Jiwoong Kim [aut, cre]

**Maintainer** Jiwoong Kim <kimjiwo2@stt.msu.edu>

**Repository** CRAN

**Date/Publication** 2015-09-14 09:12:47

## Contents

LRMDE . . . . .	1
<b>Index</b>	3

---

LRMDE	<i>Performs minimum distance estimation in linear regression model: <math>Y=Xb + \text{error}</math></i>
-------	--

---

### Description

Performs minimum distance estimation in linear regression model:  $Y=Xb + \text{error}$

### Usage

`LRMDE(Y, X)`

**Arguments**

- |   |   |
|---|---|
| Y | - Response variable in linear regression model    |
| X | - Explanatory variable in linear regression model |

**Value**

Returns betahat - Minimum distance estimator of b

**References**

- [1] Koul, H. L (1985). Minimum distance estimation in linear regression with unknown error distributions. *Statist. Probab. Lett.*, 3 1-8.
- [2] Koul, H. L (1986). Minimum distance estimation and goodness-of-fit tests in first-order autoregression. *Ann. Statist.*, 14 1194-1213.
- [3] Koul, H. L (2002). Weighted empirical process in nonlinear dynamic models. Springer, Berlin, Vol. 166

**See Also**

ARMDE

**Examples**

```
X <- matrix(c(1,1,3,4, 4,2), nrow=3, ncol=2)
Y <- c(1,-5, 8)
bhat <- LRMDE(Y,X)
```

# **Index**

LRMDE, 1